Axis I Region		Axis II System	Axis III End-organ as pain syndrome as identified from Hx, Ex and Ix	Axis IV Referral characteristics	Axis V Temporal characteristics	Axis VI Character	Axis VII Associated symptoms	Axis VIII Psychological symptoms
Chronic pelvic pain	Chronic secondary pelvic pain syndrome, formally known as specific disease associated pelvic pain OR Chronic primary pelvic pain syndrome, formally known as pelvic pain syndrome	Urological	Prostate	Suprapubic Inguinal Urethral Penile/clitoral Perineal Rectal Back Buttocks Thighs	ONSET Acute Chronic ONGOING Sporadic Cyclical Continuous TIME Filling Emptying Immediate post Late post TRIGGER Provoked Spontaneous	Aching Burning Stabbing Electric	UROLOGICAL Frequency Nocturia Hesitancy Dysfunctional flow Urgency Incontinence GYNAECOLOGICAL Menstrual Menopause GASTROINTESTINAL Constipation Diarrhoea Bloatedness Urgency Incontinence NEUROLOGICAL Dysaesthesia Hyperaesthesia Allodynia Hyperalgesia SEXUOLOGICAL Satisfaction Female dyspareunia Sexual avoidance Erectile dysfunction Medication MUSCLE Function impairment Fasciculation CUTANEOUS Trophic changes Sensory changes	ANXIETY About pain or putative cause of pain Catastrophic thinking about Pain DEPRESSION Attributed to pain or impact of pain Attributed to other causes Unattributed PTSD SYMPTOMS Re-experiencing Avoidance
			Bladder					
			Scrotal Testicular Epididymal					
			Penile Urethral					
			Post-vasectomy					
		Gynaecological	Vulvar Vestibular Clitoral					
			Endometriosis associated					
			CPPPS with cyclical exacerbations					
			Dysmenorrhoea					
		Gastrointestinal	Irritable bowel					
			Chronic anal					
			Intermittent chronic anal					
		Peripheral nerves	Pudendal pain syndrome					
		Sexological	Dyspareunia					
			Pelvic pain with sexual dysfunction					
		Psychological	Any pelvic organ					
		Musculo-skeletal	Pelvic floor muscle Abdominal muscle Spinal					
			Coccyx					

Hx = History; Ex = Examination; Ix = Investigation; PTSD = post-traumatic stress disorder.